

ORAL PRESENTATION

Open Access

# Pediatric tumours: liver tumours

Alexander J Towbin

From International Cancer Imaging Society (ICIS) 14th Annual Teaching Course Heidelberg, Germany. 9-11 October 2014

Pediatric liver tumors account for approximately 1% of all liver tumors and up to 2% of all pediatric malignancies [1,2]. While the overall incidence of liver tumors is rare, the liver remains the third most common organ of origin for solid abdominal tumors in the pediatric population [3]. The majority of pediatric liver tumors, both benign and malignant, occur (almost) exclusively in children.

Benign tumors account for one-third of all pediatric hepatic neoplasms and include infantile hepatic hemangioma, focal nodular hyperplasia, mesenchymal hamartoma, and hepatic adenoma [2-5]. Malignant neoplasms account for the remaining two-thirds of pediatric liver tumors. The most common malignancies are hepatoblastoma, hepatocellular carcinoma, fibrolamellar hepatocellular carcinoma, undifferentiated embryonal sarcoma, and biliary rhabdomyosarcoma [4-6]. While the majority of these tumors are unique to children, focal nodular hyperplasia, hepatocellular adenomas, and hepatocellular carcinoma have counterparts in adults. The presentation, risk factors, and imaging appearance of these overlapping tumors differs in the pediatric population as compared to adults [7,8].

Even though there are a number of different liver tumors, the radiologist can often make a confident diagnosis based on the clinical information and imaging appearance of the mass [9,10]. The purpose of this lecture is to describe the imaging work-up for pediatric liver masses and then discuss each tumor, focusing on its unique clinical and imaging features.

Published: 9 October 2014

## References

1. von Schweinitz D: Management of liver tumors in childhood. *Semin Pediatr Surg* 2006, **15**:17-24.
2. Hadzic N, Finegold MJ: Liver neoplasia in children. *Clin Liver Dis* 2011, **15**:443-62.

Cincinnati Children's Hospital, Department of Radiology, Cincinnati, OH 45229, USA

3. Kochin IN, Miloh TA, Arnon R, Iyer KR, Suchy FJ, Kerker N: Benign liver masses and lesions in children: 53 cases over 12 years. *Isr Med Assoc J* 2011, **13**:542-7.
4. Chung EM, Cube R, Lewis RB, Conran RM: From the archives of the AFIP: Pediatric liver masses: radiologic-pathologic correlation part 1. Benign tumors. *Radiographics* 2010, **30**:801-26.
5. Meyers RL: Tumors of the liver in children. *Surg Oncol* 2007, **16**:195-203.
6. Chung EM, Lattin GE Jr, Cube R, Lewis RB, Marichal-Hernández C, Shawhan R, Conran RM: From the archives of the AFIP: Pediatric liver masses: radiologic-pathologic correlation. Part 2. Malignant tumors. *Radiographics* 2011, **31**:483-507.
7. Towbin AJ, Luo GG, Yin H, Mo JQ: Focal nodular hyperplasia in children, adolescents, and young adults. *Pediatr Radiol* 2011, **41**:341-9.
8. Smith EA, Salisbury S, Martin R, Towbin AJ: Incidence and etiology of new liver lesions in pediatric patients previously treated for malignancy. *AJR Am J Roentgenol* 2012, **199**:186-91.
9. Meyers AB, Towbin AJ, Serai S, Geller JI, Podberesky DJ: Characterization of pediatric liver lesions with gadoxetate disodium. *Pediatr Radiol* 2011, **41**:1183-97.
10. Meyers AB, Towbin AJ, Geller JI, Podberesky DJ: Hepatoblastoma imaging with gadoxetate disodium-enhanced MRI—typical, atypical, pre- and post-treatment evaluation. *Pediatr Radiol* 2012, **42**:859-66.

doi:10.1186/1470-7330-14-S1-O16

Cite this article as: Towbin: Pediatric tumours: liver tumours. *Cancer Imaging* 2014 **14**(Suppl 1):O16.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

