

## **ORAL PRESENTATION**

**Open Access** 

## Incidental findings in the oncology patient: ovaries

Rosemarie Forstner

From International Cancer Imaging Society (ICIS) 14th Annual Teaching Course Heidelberg, Germany. 9-11 October 2014

Ovarian incidentalomas are reported in 5-18% of asymptomatic females in cross sectional imaging. Even in oncologic patients the majority of these lesions will be benign, with hydrosalpinx and ovarian cysts as leading diagnoses. Special emphasis should be given to adnexal lesions in patients with history of primaries from the GI tract or breast cancer, both of which have a propensity to metastasize to the ovaries. Patients with hereditary cancer syndromes, e. g BRCA1 or 2, are at a higher risk to develop ovarian cancer. Guidelines for the management of adnexal incidentalomas have recently been developed. Multiparametric MRI aids in further characterization of a complex adnexal lesion and thus assists in patient management in a multidisciplinary setting.

Published: 9 October 2014

doi:10.1186/1470-7330-14-S1-O46

Cite this article as: Forstner: Incidental findings in the oncology patient: ovaries. Cancer Imaging 2014 14(Suppl 1):046.

## Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit





